

GENOME REPORT

Open Access



The complete genome sequence of *Cronobacter sakazakii* ATCC 29544^T, a food-borne pathogen, isolated from a child's throat

Seongok Kim^{1,2}, You-Tae Kim³, Hyunjin Yoon², Ju-Hoon Lee^{3*†} and Sangryeol Ryu^{1†}

Abstract

Background: *Cronobacter sakazakii* is an emerging opportunistic pathogen that is associated with rare but life-threatening cases of severe diseases: meningitis, necrotizing enterocolitis, and sepsis in premature and full-term infants. However, the pathogenesis mechanism of this pathogen remains largely unknown. To determine its pathogenesis at the genomic level, the genome of *C. sakazakii* ATCC 29544^T was completely sequenced and analyzed.

Results: The genomic DNA, containing a circular chromosome and three plasmids, is composed of 4,511,265 bp with a GC content of 56.71%, containing 4380 predicted open reading frames (ORFs), 22 rRNA genes, and 83 tRNA genes. The plasmids, designated pCSK29544_p1, pCSK29544_p2, and pCSK29544_p3, were 93,905-bp, 4938-bp, and 53,457-bp with GC contents of 57.02, 54.88, and 50.07%, respectively. They were also predicted to have 72, 6, and 57 ORFs without RNA genes.

Conclusions: The strain ATCC 29544^T genome has *ompA* and *ibeB*-homologous *cusC* genes, probably associated with the invasion of human brain microvascular endothelial cells (BMECs). In addition, gene clusters for siderophore production (*iucABCD/iutA*) and the related transport system (*eitCBAD*) were detected in pCSK29544_p1 plasmid, indicating better iron uptake ability for survival. Furthermore, to survive under extremely dry condition like milk powder, this genome has gene clusters for biosynthesis of capsular proteins (CSK29544_00281-00284) and cellulose (CSK29544_01124-01127) for biofilm formation and a gene cluster for utilization of sialic acid in the milk (*nanKTAR*). The genome information of *C. sakazakii* ATCC 29544^T would provide further understanding of its pathogenesis at the molecular level for the regulation of pathogenicity and the development of a rapid detection method using biomarkers.

Keywords: *Cronobacter sakazakii*, Complete genome sequence, Pathogenesis, Infant milk formula, Virulence factor

Background

Enterobacter sakazakii has been reclassified into seven species in the genus *Cronobacter* according to biochemical and genetic evaluations [1, 2]. Among them, *Cronobacter sakazakii* is a well-known opportunistic

food-borne pathogen causing bacteremia, meningitis and necrotizing enterocolitis, particularly in low-birth-weight neonatal infants. This species is a Gram-negative, rod-shaped, peritrichous and yellow-pigmented facultative anaerobe belonging to the *Enterobacteriaceae* family [3]. *C. sakazakii* has been often found in human and infant gut microbiota [4, 5]. Although *C. sakazakii* food-borne outbreaks are quite low, the fatality to infants generally ranges from 40 to 80% [6]. Interestingly, *C. sakazakii* was reported to produce capsular or biofilm materials for its own protection from extremely dry conditions,

*Correspondence: juhlee@khu.ac.kr

[†]Ju-Hoon Lee and Sangryeol Ryu contributed equally to this work

³ Department of Food Science and Biotechnology and Institute of Life Science and Resources, Kyung Hee University, 1732 Deogyeong-daero, Giheung-gu, Yongin-si, Gyeonggi-do 17104, Republic of Korea
Full list of author information is available at the end of the article

as in formula milk powder, substantiating the high survival ability of *C. sakazakii* in the milk powder [7]. After human infection, *C. sakazakii* can invade the intestinal epithelial cells and even the brain microvascular endothelial cells (BMECs), demonstrating its potentials to cause meningitis [8]. Therefore, the biocontrol and regulation of *C. sakazakii* are urgently required. However, *C. sakazakii* is resistant to some antibiotics, indicating a problem with antibiotic therapies against *C. sakazakii* [9, 10], and its pathogenicity mechanism remains unknown. Recently, to unveil the knowledge about the *Cronobacter* ecology, multilocus sequence typing (MLST) analysis using seven housekeeping genes has been established to identify the diversity of the *Cronobacter* genus from various sources, and its application has facilitated understanding of the evolutionary relationships and environmental fitness of *Cronobacter* species [2]. In this study, to understand its infection and pathogenesis at the molecular level, the genome of a representative *C. sakazakii* type strain, ATCC 29544^T, was completely sequenced and analyzed in this study using bioinformatics. This genome information would provide the researchers with genomic insights into the virulence and pathogenicity mechanisms of this species for the further development of a rapid detection method and a novel biocontrol strategy.

Methods

Growth conditions and DNA isolation

Cronobacter sakazakii ATCC 29544^T was routinely cultivated using Luria-Bertani (LB) medium at 37 °C with shaking at 220 rpm. Bacterial cells were harvested in the mid-exponential growth phase using centrifugation at 16,000×g for 1 min and its genomic DNA was isolated using G-spinTM Genomic DNA Extraction Kit for Bacteria (iNtRON Biotechnology, Seongnam, South Korea). The concentration and purity of extracted DNA were determined by NanoVue (GE healthcare, Little Chalfont, United Kingdom).

Genome sequencing and assembly

The complete genome of *C. sakazakii* ATCC 29544^T was sequenced at Macrogen, Seoul, South Korea, using a hybrid of PacBio RS II (Pacific Biosciences, Menlo Park, CA, USA) and Illumina HiSeq 2500 (San Diego, CA, USA). The sequence reads from PacBio RS II and Illumina HiSeq 2500 platforms were assembled using HGAP (version 2.0) and ALLPATHS-LG (version r47449), respectively. The final genome coverages were on average 1321 X Illumina and 73 X PacBio, respectively.

Genome annotation

The ORFs were predicted using Glimmer3 [11] and GeneMark.hmm [12]. The gene prediction results were

confirmed by manual curation. The genes of rRNA and tRNA were predicted using RNAmmer 1.2 [13] and tRNAscan-SE [14], respectively. The genome annotation was conducted using NCBI BLASTP [15] and a predicted protein analysis using InterProScan 5 [16] for the prediction of protein functions.

The analysis of comparative genomes and phage-associated regions

Phage-associated gene clusters in the genome sequences of *C. sakazakii* ATCC 29544 were searched using PHAST server [17].

Quality assurance

Cronobacter sakazakii ATCC 29544^T was obtained from American Type Culture Collection (ATCC) and its morphological observation using a transmission electron microscopy (TEM) showed a traditional shape of *C. sakazakii* as a short rod (Additional file 1: Figure S1). In addition, this strain was confirmed to *C. sakazakii* using 16S rRNA gene sequencing. For genome sequencing, the raw read sequences were selected and assembled when their quality scores were more than 40 as cutoffs. The complete genome sequence after genome assembly was also used for confirmation using ANI analysis with previously reported complete genome sequences of *C. sakazakii*.


Initial findings

General genome properties

The complete genome of *C. sakazakii* ATCC 29544^T is composed of a circular chromosome and three plasmids (Fig. 1). The chromosome is 4,511,265 bp in DNA length with a GC content of 56.71%, 4380 ORFs, 22 rRNA genes, and 83 tRNA genes. The plasmids, designated pCSK29544_p1, pCSK29544_p2 and pCSK29544_p3, are 93,905, 4938 and 53,457 bp in DNA length with GC contents of 57.02, 54.88 and 50.07%, respectively. The plasmids have predicted ORFs of 72, 6, and 57.

Pathogenesis and virulence factors

Cronobacter sakazakii infects human neonates and infants via mostly contaminated reconstituted infant formula milk, causing serious human diseases, including bacteremia, necrotizing enterocolitis, and even meningitis [6]. The high survival rate of *C. sakazakii*, even under extremely dried conditions, as in milk formula powder, has not been investigated at the molecular level. To understand this extraordinary property of *C. sakazakii*, molecular studies have been recently performed. Interestingly, the gene clusters associated with biosynthesis of capsular polysaccharides (CSK29544_00281-00284) and cellulose (CSK29544_01124-01127) were detected. Recently, the biofilm formation and cellulose



(as a component of biofilm) production of *C. sakazakii* were experimentally confirmed [18, 19], suggesting that these gene clusters may be involved in the biofilm formation. This biofilm formation of *C. sakazakii* may contribute to the survival in the infant formula conditions [19]. However, exopolysaccharide (EPS) production was not

observed in *C. sakazakii* ATCC 29544^T [20], suggesting that capsular polysaccharide gene cluster may be associated with biofilm formation, not with EPS production. In particular, *C. sakazakii* is the only *Cronobacter* species that has the *nanKTAR* gene cluster to utilize sialic acid [21]. Interestingly, *C. sakazakii* ATCC 29544^T has this

gene cluster (CSK29544_00587-590), indicating its ability to use it. This unique ability may be involved in its adaptation to the milk conditions because sialic acid is one of the components of milk [22].

Upon human infection, *C. sakazakii* invades brain microvascular endothelial cells (BMECs), barriers to protect the brain from infection of meningitic pathogens, including *Escherichia coli* K1 and *Neisseria meningitidis*, which cause meningitis in neonates and infants [23, 24]. Outer membrane protein A (OmpA) of *C. sakazakii* is a critical determinant, contributing in vitro invasion of human BMECs by enhancing cell adhesion [24]. In addition, a few genes (*ibeA*, *ibeB*, and *yijP*) in meningitic *E. coli* were also suggested to be associated with the invasion of human BMECs [25–27]. Interestingly, two genes, *ompA* (CSK29544_03699) for BMEC adhesion and the *ibeB*-homologous *cusC* (pCSK29544_3p0028) for the penetration of BMECs, were detected in the genome of *C. sakazakii* ATCC 29544^T–, suggesting this strain may invade human BMECs. However, *ibeA* and *yijP* were not detected in the genome. It is noteworthy that this *ibeB*-homologous *cusC* is located in the gene cluster of the complete copper- and silver-resistance cation efflux system as encoded by *cusABCF* (CSK29544_3p0026–CSK29544_3p0031) in the plasmid pCSK29544_3p, suggesting this BMEC invasion ability may be transferrable to other *C. sakazakii* or that the strain ATCC 29544^T may have been acquired from other *C. sakazakii* for human BMEC invasion. The presence of a conjugational transfer protein encoded by *traX* (CSK29544_3p0007) supports this hypothesis.

Cronobacter sakazakii ATCC 29544^T harbors three plasmids, pCSK29544_1p (1p), pCSK29544_2p (2p), and pCSK29544_3p (3p). Interestingly, two large plasmids (1p and 3p) encode virulence-associated genes, related to iron uptake and BMEC invasion. Therefore, these two plasmids may contribute to host dominance and pathogenesis, respectively.

Iron is an essential element for dominant survival and colonization via bacterial competition for iron uptake because it plays an important role in the electron transport system for energy production [28]. To accomplish this dominant survival and colonization against other bacteria, *C. sakazakii* has an iron acquisition system, including siderophore production (*iucABCD/iutA*) and an ABC-type transport system (*eitCBAD*) in the plasmid pESA3 [29]. The strain ATCC 29544^T also has this privileged iron acquisition system, including siderophore biosynthesis (*iucABCD/iutA*; CSK29544_1p0024–CSK29544_1p0028) and an ABC-type iron transport system (*eitCBAD*; CSK29544_1p0056–CSK29544_1p0059). However, this iron acquisition system is present in the plasmid 1p, similar to pESA3 of *C. sakazakii* BAA-894 [29]. Interestingly,


C. sakazakii BAA-894 has multiple copies of pESA3 plasmids in a host, which is highly homologous to the plasmid 1p, suggesting the plasmid 1p may be multi copied, too [30]. These results indicate that the host strain may take advantage of better iron uptake ability in the given environments, probably due to the presence of multiple copies of this iron acquisition system encoded in the plasmid 1p.

Recent studies have revealed that milk formula contains an at least six-times-higher arsenic concentration than that of breast milk, suggesting bacterial survival in milk formula may require arsenic resistance [31]. In addition to the previously mentioned BMEC invasion ability of *C. sakazakii* ATCC 29544^T via the *ibeB*-homologous *cusC* gene in the cation efflux system, the plasmid 3p has an arsenic resistance system as encoded by *arsRDABC*, suggesting the strain ATCC 29544^T may have arsenic resistance activity for survival in formula milk powder and that this ability may have been acquired via plasmid conjugational transfer.

Comparative genome analysis

To date, five complete genome sequences of *C. sakazakii* and their annotation data are available in the GenBank database. To compare these genomes, whole genomic DNA sequence-based average nucleotide index (ANI) analysis and Roary matrix-based protein sequence analysis were performed. ANI analysis showed that *C. sakazakii* NCTC 8155 and SP291 are the most closely related, forming a taxonomical group, and the strains ATCC 29544^T, ATCC BAA894, and ES15 are also similar to this group (Fig. 2a). Subsequent Roary matrix-based protein sequence analysis of these five genomes supports this relationship among them (Fig. 2b). However, the strain ATCC 29544^T is more closely related to the group in the Roary matrix-based tree.

Moreover, pan genome analysis of these five genomes was conducted and their strain-specific regions were detected (Fig. 1). Interestingly, three functional gene clusters and seven prophage regions are ATCC 29544^T-specific. In addition to the iron acquisition system consisting of siderophore production and ABC-type transport system in the plasmid 1p, ATCC 29544^T-specific iron uptake-associated gene cluster containing *fecABDCEI* for dominant uptake of ferric dicitrate is present in the chromosome, probably related to survival in the iron-limited condition [32]. The second arsenic resistance gene cluster consisting of *arsRCB* for arsenate sensing, reduction, and efflux pumping was also detected, which is similar to the gene cluster in the plasmid 3p, probably for dominant survival in the human gut containing low concentration of arsenic after human infection [33] (Additional file 1: Fig. S2A). Comparative analysis of their protein sequences showed that they are >94% identical



to each other, probably for synergistic arsenic resistance activity. Unlike other two gene clusters shared by the chromosome and plasmid, two copies of lactose operons are present in the chromosomes of ATCC 29544^T, NCTC 8155, and SP291, probably related to the Roary matrix-based tree (Fig. 2b). While one of these operons (CSK29544_04271–04273) are shared by all *C. sakazakii* genomes, the second lactose operon (CSK29544_00439–00442) is ATCC 29544^T-specific. This second lactose operon is similar to other lactose operon in *Klebsiella* and *Enterobacter* (data not shown), suggesting that this operon may be originated from other genus bacterium. The presence of multiple mobile elements containing transposases and relatively low protein sequence identity support this hypothesis (Additional file 1: Fig. S2B). Furthermore, pan genome analysis showed strain-specific region (Fig. 1). Most of genes in this region belong to prophages and they have very low homology among them, indicating that these prophages are also strain-specific. Interestingly, the strain ATCC 29544^T contains seven strain-specific prophage regions with five intact prophages and two incomplete prophages (Table 1). However, most of genes in the prophages encode hypothetical proteins.

Future directions

This genome information of *C. sakazakii* ATCC 29544^T provides genomic insights into the presence of various virulence factors and their associated pathogenesis mechanisms at molecular level. In addition, this genome information showed that the strain ATCC 29544^T has

Table 1 Genome statistics

Attribute	Value	% of Total ^a
Genome size (bp)	4,663,565	100
DNA coding (bp)	4,116,350	88.27
DNA G + C (bp)	2,641,511	56.64
Total genes	4620	100
Protein coding genes	4515	97.72
RNA genes	105	2.27
rRNA operons	22	0.48
tRNA genes	83	1.8
Genes with function prediction	3212	69.52
Predicted prophages	7	6.18
CRISPR repeats	2	0.04

^a The total is based on either the size of the genome (chromosome and 3 plasmids) in base pairs or the total number of total genes in the annotated genome

several distinct features probably contributing to dominant survival under the given habitats and even various stress conditions like extremely dry or nutrient-limited conditions. Therefore, this genome information would be useful for further development of a novel strategy for control of *C. sakazakii* and of rapid detection method using strain-specific DNA markers in genomic level.

Additional file

Additional file 1: Figure S1. The morphology of *C. sakazakii* ATCC 29544 imaged by energy-filtering transmission electron microscopy (EF-TEM). EF-TEM photograph was obtained by 2% uranyl acetate on copper grids and examined with ET-TEM at a voltage of 120 kV (LIBRA 120, Zeiss, Oberkochen, Germany). **Figure S2:** Comparative analysis of two ATCC 29544-specific gene clusters: (A) *lac* operon and (B) arsenic resistance. The amino acid sequence identities between associated genes are indicated as percentages.

Abbreviations

BMECs: brain microvascular endothelial cells; ANI: average nucleotide identity; LB: Luria–Bertani medium; MLST: multilocus sequence typing; TEM: transmission electron microscopy; COG: clusters of orthologous groups.

Authors' contributions

SK, JHL, and SR initiated and supervised the study. SK and JHL drafted the manuscript. SK conducted laboratory experiments, and SK and YTK worked on the genome sequencing and annotated the genome. SK, YTK, HY, JHL, and SR discussed and analyzed the data and revised the manuscript. All authors read and approved the final manuscript.

Author details

¹ Department of Food and Animal Biotechnology, Department of Agricultural Biotechnology, Research Institute for Agriculture and Life Sciences, and Center for Food and Bioconvergence, Seoul National University, 1 Gwanak-ro, Gwanak-gu, Seoul 08826, Republic of Korea. ² Department of Applied Chemistry and Biological Engineering, Ajou University, 206 Worldcup-ro, Yeongtong-gu, Suwon 16499, Republic of Korea. ³ Department of Food Science and Biotechnology and Institute of Life Science and Resources, Kyung Hee University, 1732 Deogyeong-daero, Giheung-gu, Yongin-si, Gyeonggi-do 17104, Republic of Korea.

Acknowledgements

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

The genome sequences of *C. sakazakii* ATCC 29544^T were deposited in the NCBI GenBank server under the accession numbers CP011047, CP011048, CP011049, and CP011050 for a chromosome and three plasmids, respectively.

Funding

This research was supported by a grant (14162MFDS972) from the Ministry of Food and Drug Safety, Korea, in 2016 and by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT and Future Planning (NRF-2015R1C1A1A01053815).

Received: 11 November 2016 Accepted: 7 December 2016

Published online: 04 January 2017

References

- Iversen C, Lehner A, Mullane N, Bidlas E, Cleenwerck I, Marugg J, et al. The taxonomy of *Enterobacter sakazakii*: proposal of a new genus *Cronobacter*

- gen. nov. and descriptions of *Cronobacter sakazakii* comb. nov. *Cronobacter sakazakii* subsp. *sakazakii*, comb. nov., *Cronobacter sakazakii* subsp. *malonaticus* subsp. nov., *Cronobacter turicensis* sp. nov., *Cronobacter muytjensii* sp. nov., *Cronobacter dublinensis* sp. nov. and *Cronobacter genomospecies* 1. BMC Evol Biol. 2007;7:64.
- Joseph S, Forsythe SJ. Insights into the emergent bacterial pathogen *Cronobacter* spp., generated by multilocus sequence typing and analysis. Front Microbiol. 2012;3:397.
- Farmer JJ, Asbury MA, Hickman FW, Brenner DJ. *Enterobacter sakazakii*—a new species of *Enterobacteriaceae* isolated from clinical specimens. Int J Syst Bacteriol. 1980;30:569–84.
- Hallstrom M, Eerola E, Vuento R, Janas M, Tammela O. Effects of mode of delivery and necrotising enterocolitis on the intestinal microflora in preterm infants. Eur J Clin Microbiol Infect Dis. 2004;23:463–70.
- Mao B, Li D, Zhao J, Liu X, Gu Z, Chen YQ, et al. In vitro fermentation of fructooligosaccharides with human gut bacteria. Food Funct. 2015;6:947–54.
- Healy B, Cooney S, O'Brien S, Iversen C, Whyte P, Nally J, et al. *Cronobacter* (*Enterobacter sakazakii*): an opportunistic foodborne pathogen. Food-borne Pathog Dis. 2010;7:339–50.
- Hurrell E, Kucerova E, Loughlin M, Caubilla-Barron J, Forsythe SJ. Biofilm formation on enteral feeding tubes by *Cronobacter sakazakii*, *Salmonella* serovars and other *Enterobacteriaceae*. Int J Food Microbiol. 2009;136:227–31.
- Giri CP, Shima K, Tall BD, Curtis S, Sathyamoorthy V, Hanisch B, et al. *Cronobacter* spp. (previously *Enterobacter sakazakii*) invade and translocate across both cultured human intestinal epithelial cells and human brain microvascular endothelial cells. Microb Pathog. 2012;52:140–7.
- Kilonzo-Nthenge A, Rotich E, Godwin S, Nahashon S, Chen F. Prevalence and antimicrobial resistance of *Cronobacter sakazakii* isolated from domestic kitchens in middle Tennessee, United States. J Food Prot. 2012;75:1512–7.
- Xu X, Li C, Wu Q, Zhang J, Huang J, Yang G. Prevalence, molecular characterization, and antibiotic susceptibility of *Cronobacter* spp. in Chinese ready-to-eat foods. Int J Food Microbiol. 2015;204:17–23.
- Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. Improved microbial gene identification with GLIMMER. Nucleic Acids Res. 1999;27:4636–41.
- Lukashin AV, Borodovsky M. GeneMark.hmm: new solutions for gene finding. Nucleic Acids Res. 1998;26:1107–15.
- Lagesen K, Hallin P, Rodland EA, Staerfeldt HH, Rognes T, Ussery DW. RNAMmer: consistent and rapid annotation of ribosomal RNA genes. Nucleic Acids Res. 2007;35:3100–8.
- Lowe TM, Eddy SR. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. Nucleic Acids Res. 1997;25:955–64.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990;215:403–10.
- Jones P, Binns D, Chang HY, Fraser M, Li W, McAnulla C, et al. InterPro-Scan 5: genome-scale protein function classification. Bioinformatics. 2014;30:1236–40.
- Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster version of the PHAST phage search tool. Nucleic Acids Res. 2016;44:W16–21.
- Choi Y, Kim S, Hwang H, Kim KP, Kang DH, Ryu S. Plasmid-encoded MCP is involved in virulence, motility, and biofilm formation of *Cronobacter sakazakii* ATCC 29544. Infect Immun. 2015;83:197–204.
- Hu L, Grim CJ, Franco AA, Jarvis KG, Sathyamoorthy V, Kothary MH, et al. Analysis of the cellulose synthase operon genes, *bcsA*, *bcsB*, and *bcsC* in *Cronobacter* species: prevalence among species and their roles in biofilm formation and cell-cell aggregation. Food Microbiol. 2015;52:97–105.
- Lehner A, Riedel K, Eberl L, Breeuwer P, Diep B, Stephan R. Biofilm formation, extracellular polysaccharide production, and cell-to-cell signaling in various *Enterobacter sakazakii* strains: aspects promoting environmental persistence. J Food Prot. 2005;68:2287–94.
- Joseph S, Hariri S, Masood N, Forsythe S. Sialic acid utilization by *Cronobacter sakazakii*. Microb Inform Exp. 2013;3:3.
- Wang B, Brand-Miller J, McVeagh P, Petocz P. Concentration and distribution of sialic acid in human milk and infant formulas. Am J Clin Nutr. 2001;74:510–5.
- Nassif X, Bourdoulous S, Eugene E, Couraud PO. How do extracellular pathogens cross the blood-brain barrier? Trends Microbiol. 2002;10:227–32.

24. Nair MK, Venkitanarayanan K, Silbart LK, Kim KS. Outer membrane protein A (OmpA) of *Cronobacter sakazakii* binds fibronectin and contributes to invasion of human brain microvascular endothelial cells. *Foodborne Pathog Dis*. 2009;6:495–501.
25. Huang SH, Wan ZS, Chen YH, Jong AY, Kim KS. Further characterization of *Escherichia coli* brain microvascular endothelial cell invasion gene *ibeA* by deletion, complementation, and protein expression. *J Infect Dis*. 2001;183:1071–8.
26. Huang SH, Chen YH, Fu Q, Stins M, Wang Y, Wass C, et al. Identification and characterization of an *Escherichia coli* invasion gene locus, *ibeB*, required for penetration of brain microvascular endothelial cells. *Infect Immun*. 1999;67:2103–9.
27. Wang Y, Huang SH, Wass CA, Stins MF, Kim KS. The gene locus *yijP* contributes to *Escherichia coli* K1 invasion of brain microvascular endothelial cells. *Infect Immun*. 1999;67:4751–6.
28. Mietzner TA, Morse SA. The role of iron-binding proteins in the survival of pathogenic bacteria. *Annu Rev Nutr*. 1994;14:471–93.
29. Franco AA, Hu L, Grim CJ, Gopinath G, Sathiamoorthy V, Jarvis KG, et al. Characterization of putative virulence genes on the related RepFIB plasmids harbored by *Cronobacter* spp. *Appl Environ Microbiol*. 2011;77:3255–67.
30. Kucerova E, Clifton SW, Xia XQ, Long F, Porwollik S, Fulton L, et al. Genome sequence of *Cronobacter sakazakii* BAA-894 and comparative genomic hybridization analysis with other *Cronobacter* species. *PLoS ONE*. 2010;5:e9556.
31. Carignan CC, Cottingham KL, Jackson BP, Farzan SF, Gandolfi AJ, Punshon T, et al. Estimated exposure to arsenic in breastfed and formula-fed infants in a United States cohort. *Environ Health Perspect*. 2015;123:500–6.
32. Harvie DR, Ellar DJ. A ferric dicitrate uptake system is required for the full virulence of *Bacillus cereus*. *Curr Microbiol*. 2005;50:246–50.
33. Ratnaik RN. Acute and chronic arsenic toxicity. *Postgrad Med J*. 2003;79:391–6.
34. Petkau A, Stuart-Edwards M, Stothard P, Van Domselaar G. Interactive microbial genome visualization with GView. *Bioinformatics*. 2010;26:3125–6.
35. Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, et al. Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics*. 2015;31:3691–3.

Submit your next manuscript to BioMed Central
and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

